arXiv:math/0611080v2 [math.SG] 29 Sep 2008

LEGENDRIAN HELIX AND CABLE LINKS

FAN DING AND HANSJÖRG GEIGES

Abstract. Lisa Traynor has described an example of a two-component Legendrian 'circular helix link' $\Lambda_0 \sqcup \Lambda_1$ in the 1–jet space $J^1(S^1)$ of the circle (with its canonical contact structure) that is topologically but not Legendrian isotopic to the link $\Lambda_1 \sqcup \Lambda_0$. We give a complete classification of the Legendrian realisations of this topological link type, as well as all other 'cable links' in $J^1(S^1)$.

1. Introduction

Considerable progress has been made towards the classification of Legendrian knots and links in contact 3–manifolds, e.g. [3, 5, 9, 1]. For a general introduction to this topic see [2]. In [10], Lisa Traynor exhibited an intriguing example of an ordered Legendrian two-component link that is topologically but not Legendrian isotopic to the link obtained by interchanging the two components. The purpose of the present paper is to give a complete classification of the Legendrian realisations of this and some related topological link types.

Let

$$J^1(S^1) = T^*S^1 \times \mathbb{R} = \{(x, y, z)\colon\ x \in \mathbb{R}/2\pi\mathbb{Z},\ y, z \in \mathbb{R}\}$$

be the 1–jet space of S^1 with its standard contact structure

$$\xi = \ker(dz - y\,dx).$$

Here y denotes the fibre coordinate in T^*S^1, and z the coordinate in the $\mathbb{R}$–factor of $J^1(S^1)$. The graph of any smooth function $g\colon\ S^1 \to \mathbb{R}$ lifts to a Legendrian knot

$$\Lambda_g := \{(x, y, z)\colon\ y = g'(x),\ z = g(x)\}.$$

The linear interpolation between two smooth functions $g, h\colon\ S^1 \to \mathbb{R}$ gives rise to a Legendrian isotopy between Λ_g and Λ_h. The following theorem, therefore, is rather surprising. Here we write Λ_s for the Legendrian knot corresponding to the constant function of value $s \in \mathbb{R}$, oriented by the variable x.

Theorem 1 (Traynor)**.** *The ordered Legendrian links $\Lambda_0 \sqcup \Lambda_1$ and $\Lambda_1 \sqcup \Lambda_0$ in $(J^1(S^1), \xi)$ are topologically, but not Legendrian isotopic.*

In Theorem 2 below we give a complete classification of the Legendrian realisations of the 'circular helix link', i.e. the topological link type $\Lambda_0 \sqcup \Lambda_1$ in $(J^1(S^1), \xi)$. We then take this theorem as the starting point for the classification of more general Legendrian 'cable links' in $(J^1(S^1), \xi)$. A further purpose of the present paper is to address a subtle point in our earlier classification of Legendrian cable links in S^3, where previously we did not provide full details, see [1, p. 154]. The relevant argument relies in an essential way on the classification of Legendrian circular helix links.

Before stating our results, we need to recall the definition of the classical invariants of a Legendrian knot in $(J^1(S^1),\xi)$.

2. Legendrian knots in $J^1(S^1)$

By identifying S^1 with $\mathbb{R}/2\pi\mathbb{Z}$, we can visualise a Legendrian knot $K\subset J^1(S^1)$ in its front projection to a strip $[0,2\pi]\times\mathbb{R}$ in the xz–plane. The **Thurston–Bennequin invariant** of K is

$$\mathtt{tb}(K)=\mathrm{writhe}(K)-\frac{1}{2}\#(\mathrm{cusps}(K)),$$

where the quantities on the right are computed from the front projection of K. This is the signed number of crossing changes required to 'unlink' K from its push-off K' in the z–direction (i.e. transverse to ξ), that is, the number of crossing changes that will allow one to separate the two knots in $J^1(S^1)$.

The **rotation number** of the *oriented* Legendrian knot K is defined in terms of the front projection as

$$\mathtt{rot}(K)=\frac{1}{2}(c_- - c_+),$$

with $c_\pm$ the number of cusps oriented upwards or downwards, respectively.

There are definitions of these invariants that do not rely on the front projection, and which show that $\mathtt{tb}$ and $\mathtt{rot}$ are in fact Legendrian isotopy invariants, cf. [1].

The discussion in [1] also shows that the Thurston–Bennequin inequality for Legendrian realisations L_0 of the topological knot type Λ_0 takes the form

$$\mathtt{tb}(L_0)+|\mathtt{rot}(L_0)|\leq 0.$$

It follows that the maximal Thurston–Bennequin invariant of such knots L_0 is $\overline{\mathtt{tb}}(L_0)=0$, and this maximal value is realised by the Legendrian knot Λ_0. By stabilising Λ_0, as described in the next section, one can realise any non-positive integer $-m$ in the form $-m=\mathtt{tb}(L_0)$.

In [1, Section 6] we wrote down an explicit contact embedding $f\colon J^1(S^1)\to S^3$ (both manifolds equipped with their standard contact structure). The image of f is the complement of a standard Legendrian unknot K_0; the knots $f(\Lambda_0)$ and K_0 form a Hopf link. We shall frequently use the contactomorphism $f\colon J^1(S^1)\to S^3\setminus K_0$ to relate Legendrian links in the two manifolds.

3. Legendrian helix links

The **stabilisation** $S_\pm(K)$ of a Legendrian knot K in $(J^1(S^1),\xi)$ is the Legendrian knot whose front projection is obtained from that of K by adding a 'zigzag' as in Figure 1.

Figure 1. Legendrian 'zigzags'.

A zigzag oriented downwards gives a positive stabilisation S_+, while a zigzag oriented upwards gives S_-. Hence

$$\mathtt{tb}(S_\pm(K))=\mathtt{tb}(K)-1,\ \ \mathtt{rot}(S_\pm(K))=\mathtt{rot}(K)\pm 1.$$

Stabilisations are well defined and commute with each other.

Theorem 2. *Let $L = L_0 \sqcup L_1$ be a Legendrian link in $(J^1(S^1), \xi)$ of the same (topological) oriented link type as $\Lambda_0 \sqcup \Lambda_1$.*

(a) If $\mathtt{tb}(L_0) = \mathtt{tb}(L_1) = 0$, then L is Legendrian isotopic to either $\Lambda_0 \sqcup \Lambda_1$ or $\Lambda_1 \sqcup \Lambda_0$.

(b) If one of $\mathtt{tb}(L_0), \mathtt{tb}(L_1)$ is negative, then L is Legendrian isotopic to

$$S_+^{k_0} S_-^{l_0}(\Lambda_0) \sqcup S_+^{k_1} S_-^{l_1}(\Lambda_1),$$

where

$$k_i = \frac{-\mathtt{tb}(L_i) + \mathtt{rot}(L_i)}{2}, \quad l_i = \frac{-\mathtt{tb}(L_i) - \mathtt{rot}(L_i)}{2}, \quad i = 0, 1.$$

In particular, the phenomenon observed by Traynor disappears with the first stabilisation. Theorem 2 will be proved by a method analogous to that in [3, 1]. In [1], one of the key ingredients was the classification of tight contact structures on a thickened torus $T^2 \times [0, 1]$, here it is the classification of tight contact structures on the product of a pair of pants with a circle, as obtained by Etnyre and Honda [4]. The other main ingredient is convex surface theory in the sense of Giroux; the exposition given in [3, Section 3] is sufficient for our purposes.

Proof. (a) For $R > 0$, consider the solid torus

$$M_R = \{(x, y, z) \in J^1(S^1) \colon\ y^2 + z^2 \leq R^2\}.$$

As meridian μ of this solid torus we may take the curve on ∂M_R given by $x = 0$, oriented positively in the yz–plane; the curve given by $(y, z) = (R, 0)$, oriented by the parameter x, can serve as longitude λ. Then (μ, λ) is a positive basis for $H_1(\partial M_R)$.

The vector field $X := y\partial_y + z\partial_z$ satisfies $\mathcal{L}_X(dz - y\,dx) = dz - y\,dx$, so it is a contact vector field for ξ. Hence ∂M_R is a convex torus whose dividing set — the set of points where $X \in \xi$ — consists of two longitudes $\{y = \pm R, z = 0\}$; i.e. these dividing curves have slope ∞ with respect to (μ, λ). The characteristic foliation on ∂M_R is the Legendrian ruling of slope 1 defined by the vector field $\partial_x + y\partial_z - z\partial_y$; this foliation has singularities along the Legendrian divides $\{y = 0, z = \pm R\}$.

Since $\mathtt{tb}(\Lambda_0) = \mathtt{tb}(\Lambda_1) = 0$, we find disjoint tubular neighbourhoods N_0, N_1 of these two knots with convex boundary having two dividing curves of slope ∞. Likewise, we may choose disjoint tubular neighbourhoods V_0, V_1 of L_0, L_1, respectively, with convex boundary of the same kind. These neighbourhoods are contactomorphic by a diffeomorphism preserving the longitude (and, obviously, the meridian). We may assume that R has been chosen large enough such that these four tubular neighbourhoods are contained in $\mathrm{Int}(M_R)$.

Write Σ for a pair of pants, i.e. a disc with two open discs removed. Both $M_R \setminus \mathrm{Int}(N_0 \cup N_1)$ and $M_R \setminus \mathrm{Int}(V_0 \cup V_1)$ are homeomorphic to $\Sigma \times S^1$. All the boundary tori are convex of slope ∞ with respect to ξ. As shown in Lemma 10 and Lemma 11 of [4], this information determines the tight contact structure on $\Sigma \times S^1$ up to a permutation of the two interior 'holes'; in other words, we find a contactomorphism

$$\phi \colon\ M_R \setminus \mathrm{Int}(V_0 \cup V_1) \longrightarrow M_R \setminus \mathrm{Int}(N_0 \cup N_1)$$

that is the identity on ∂M_R, and sends $(\partial V_0, \partial V_1)$ to $(\partial N_{s_0}, \partial N_{s_1})$ for some permutation (s_0, s_1) of $(0, 1)$, preserving the corresponding longitudes and meridians.

We can extend ϕ to a contactomorphism of $(J^1(S^1),\xi)$ that sends $L_0 \sqcup L_1$ to $\Lambda_{s_0} \sqcup \Lambda_{s_1}$ and equals the identity map outside M_R. As explained in the proof of [1, Thm. 3.3], such a contactomorphism is contact isotopic to the identity. This yields the Legendrian isotopy of $L_0 \sqcup L_1$ to $\Lambda_{s_0} \sqcup \Lambda_{s_1}$.

(b) Suppose that $\mathtt{tb}(L_0) < 0$. Choose an annulus A, disjoint from L_1, with boundary curves L_0 and Λ_s for some large constant s. The surface framing of Λ_s determined by A corresponds to the vector field ∂_z, so does the contact framing of Λ_s, which satisfies $\mathtt{tb}(\Lambda_s) = 0$. Since $\mathtt{tb}(L_0) < 0$, the contact framing of L_0 determined by ξ makes at least one negative (i.e. left) twist relative to its surface framing determined by A. The condition that the contact structure ξ twist non-positively relative to the surface framing along either boundary component of A is precisely what is needed so that one may invoke [6, Prop. 3.1], which tells us that we can find a C^0–small perturbation of A rel L_0, Λ_s (keeping it disjoint from L_1) into a convex surface.

In slightly less explicit form, this perturbation statement is also part of the following result of Kanda [7], cf. [1, Thm. 4.8], to which we shall refer frequently in the sequel.

Theorem 3 (Kanda)**.** *If γ is a Legendrian curve in a surface Σ, then Σ may be isotoped relative to γ so that it is convex if and only if the twisting $t_\Sigma(\gamma)$ of the contact planes along γ relative to the framing induced by Σ satisfies $t_\Sigma(\gamma) \leq 0$. If Σ is convex, then*

$$t_\Sigma(\gamma) = -\frac{1}{2}\#(\gamma \cap \Gamma),$$

where $\#(\gamma \cap \Gamma)$ denotes the number of intersection points of γ with the dividing set Γ of Σ.

By this theorem, the dividing curves of the (convex) annulus A intersect L_0, Λ_s in $-2\mathtt{tb}(L_0) > 0$, $-2\mathtt{tb}(\Lambda_s) = 0$ points, respectively. This means that there is a boundary parallel dividing curve along L_0, which allows us to destabilise this Legendrian knot in $J^1(S^1) \setminus L_1$, cf. [3, Lemma 3.9].

By iterating this procedure we find Legendrian knots L_0', L_1' with $\mathtt{tb}(L_0') = \mathtt{tb}(L_1') = 0$ and non-negative integers k_i, l_i such that $S_+^{k_i} S_-^{l_i}(L_i') = L_i$, $i = 0, 1$. So, by (a), L is Legendrian isotopic to

$$S_+^{k_0} S_-^{l_0}(\Lambda_{s_0}) \sqcup S_+^{k_1} S_-^{l_1}(\Lambda_{s_1})$$

for some permutation (s_0, s_1) of $(0, 1)$. From the behaviour of the classical invariants $\mathtt{tb}$ and $\mathtt{rot}$ under stabilisation it follows that the relation between k_i, l_i and $\mathtt{tb}(L_i), \mathtt{rot}(L_i)$ is as stated in the theorem.



FIGURE 2. A Legendrian isotopy between $\Lambda_0 \sqcup S_+(\Lambda_1)$ and $\Lambda_1 \sqcup S_+(\Lambda_0)$.

It remains to show that in fact either permutation gives the same Legendrian isotopy class, provided there is at least one stabilisation. For that, it suffices to

describe a Legendrian isotopy between, say, $\Lambda_0 \sqcup S_+(\Lambda_1)$ and $\Lambda_1 \sqcup S_+(\Lambda_0)$; the picture for a single negative stabilisation is analogous. Figure 2 indicates that a couple of Legendrian Reidemeister moves of the second kind will do the job. □

4. Cable links in S^3 revisited

In our earlier paper [1] we gave a classification of Legendrian (p, q)–cable links in S^3. As pointed out on p. 154 of that paper, we still owe the reader a final detail in one particular case of that classification. Here we provide the missing argument, which depends crucially on Theorem 2.

Recall the set-up from [1], cf. Figure 3. Let L_0 be a trivial knot in S^3. Denote by μ and λ meridian and longitude on a torus T, viewed as the boundary of the *complement* of a tubular neighbourhood of L_0; this unusual convention was chosen in [1] for technical reasons. A (p, q)–**cable link** is a link of the form $\mathbb{L} = L_0 \sqcup L_1$ with L_1 a knot on T homologically equivalent to $p\mu + q\lambda$.



Figure 3. Choice of meridian μ and longitude λ.

In [1] we classified Legendrian realisations of this link type. With the notation $m := -\mathtt{tb}(L_0)$, the case for which additional details have to be provided is the one where $q \geq 2$, $p = -1$ and $-m + q \leq 0$ (case 3(b2-iii) in [1]). Here the maximal possible value of $\mathtt{tb}(L_1)$ is pq. What remains to be shown in order to complete the argument from [1] is the uniqueness of the Legendrian realisation of this link type with $\mathtt{tb}(L_1) = pq$ and given rotation numbers in the allowable range determined by equations (4.3) and (4.4) of [1] (as we shall see, no explicit reference to these equations will be necessary in the discussion that follows).

As explained in [1, p. 147], we may assume that the torus T on which L_1 sits is a convex torus in standard form (cf. [1, Defn. 4.1]), and L_1 is one of an even number of Legendrian divides on T. The slope of these curves — with T now regarded as the boundary of a tubular neighbourhood of L_0 — is $p/q = -1/q$.

For such slopes, [3, Lemma 3.17] does not apply. For other slopes, this lemma could be used to show that all Legendrian divides are isotopic. In particular, one could use bypasses in order to remove all but two Legendrian divides and still assume without loss of generality that L_1 was one of these remaining Legendrian divides.

Instead, we argue as follows. The torus T being convex, there is a contact flow transverse to T. Use this flow to push T further away from L_0, so that $\mathbb{L}$ may now be regarded as a link in the interior of a solid torus V with convex boundary of slope $-1/q$. If there are more than two dividing curves (and hence Legendrian

divides) on ∂V we can use bypasses on meridional discs of the complementary solid torus for reducing the number of dividing curves down to 2. Now ignoring the complement of V, we can apply a diffeomorphism to V (changing the longitude(!) of ∂V from $-\mu$ to $-\mu+q\lambda$) such that the boundary slope becomes ∞.

This allows us to identify V with a standard neighbourhood of Λ_0 in $J^1(S^1)$. With respect to this identification, $\mathbb{L}$ is topologically isotopic to $\Lambda_0 \sqcup \Lambda_1$. If $-m+q<0$ then we are done by Theorem 2 (b), because in this situation Legendrian realisations of this link are unique even inside V.

If $-m+q=0$, then by Theorem 2 (a) our link $\mathbb{L}$ is Legendrian isotopic inside V to one of $\Lambda_0 \sqcup \Lambda_1$ or $\Lambda_1 \sqcup \Lambda_0$. Observe that the two components of this link are Legendrian push-offs of each other. Translated back into S^3, this means that $\mathbb{L}$ consists of a topologically trivial Legendrian knot and its Legendrian push-off, or what is called the 2–**copy** of a topologically trivial Legendrian knot in [8]. (For the orientations to be consistent with the interpretation of L_1 as the push-off of L_0, however, we need to replace L_1 by its reverse $\overline{L}_1$.) As a consistency check, observe that $\mathtt{tb}(L_0)=-m=-q=pq=\mathtt{tb}(\overline{L}_1)$ and $\mathtt{lk}(L_0,\overline{L}_1)=-q=\mathtt{tb}(L_0)$. Moreover, formulæ (4.3) and (4.4) in [1] give $\mathtt{rot}(L_0)=\mathtt{rot}(\overline{L}_1)$.

As shown in [8, Prop. 4.2a,b], cyclic permutations (by a Legendrian isotopy) of the N–copy of a topologically trivial Legendrian knot are possible in S^3 (in contrast with Theorem 2 (a)). This proves the uniqueness of the Legendrian realisations of $\mathbb{L}$ in this case and completes the argument from [1].

Remark. According to [8, Thm. 5.1a], only cyclic permutations of the N–copy of the Legendrian unknot are possible. For $N=3$, this result is equivalent to Theorem 1. Indeed, with $f\colon J^1(S^1)\to S^3\setminus K_0$ being the contactomorphism mentioned earlier, the Legendrian link $f(\Lambda_0)\sqcup f(\Lambda_1)\sqcup K_0$ is the 3–copy of the Legendrian unknot. Thus, if one were able to permute Λ_0 and Λ_1 in $J^1(S^1)$, one would have a transposition of $f(\Lambda_0)$ and $f(\Lambda_1)$, even with K_0 fixed during the isotopy. Conversely, if one could transpose $f(\Lambda_0)$ and $f(\Lambda_1)$ by a Legendrian isotopy ϕ_t that ends up moving K_0 to itself, one could also find a contactomorphism of S^3 that transposes $f(\Lambda_0)$ and $f(\Lambda_1)$, and fixes a neighbourhood of K_0 pointwise. (Define the contactomorphism to be the time–1–map ϕ_1 of the isotopy outside a standard tubular neighbourhood N_0 of K_0, set it equal to the identity on a smaller standard tubular neighbourhood contained in the interior of both N_0 and $\phi_1(N_0)$, and extend this to a contactomorphism on all of S^3 using the uniqueness up to contactomorphism of the non-rotative tight contact structure on $T^2\times[0,1]$ with two dividing curves on each boundary component, as proved in [6, Prop. 4.9].) Our argument in [1, Proof of Thm. 3.3] then shows that one could also find a Legendrian isotopy that transposes $f(\Lambda_0)$ and $f(\Lambda_1)$ and keeps a neighbourhood of K_0 fixed during the isotopy. Such an isotopy would induce a permutation of Λ_0 and Λ_1 in $J^1(S^1)$.

5. Cable links in $J^1(S^1)$

Recall the definition of M_R and μ,λ given at the beginning of the proof of Theorem 2. Any torus of the form $T=\partial M_R$ for some $R>0$ is called a **standard torus**. By a (p,q)–**cable link** in $J^1(S^1)$ we mean a link $\mathbb{L}=L_0\sqcup L_1$ isotopic (as an ordered, oriented link) to the union of Λ_0 and a (p,q)–cable of Λ_0, i.e. a knot on a standard torus in the class of $p\mu+q\lambda$ (with p and q coprime). Our main theorem gives a classification of the Legendrian realisations of this link type.

Theorem 4. *With the exception of the case in Theorem 2 (a), two oriented Legendrian cable links in $(J^1(S^1), \xi)$ are Legendrian isotopic if and only if their oriented link types and their classical invariants* `tb` *and* `rot` *agree.*

For a given pair (p, q), let $\mathbb{L} = L_0 \sqcup L_1$ and $\mathbb{L}^* = L_0^* \sqcup L_1^*$ be two Legendrian realisations of the (p, q)–cable link having the same classical invariants. According to [1, Thm. 3.3], Legendrian torus knots in $(J^1(S^1), \xi)$ are determined by the classical invariants. Applied to L_0 and L_0^*, regarded as a $(0, 1)$–torus knot, this result tells us that we may assume $L_0 = L_0^* = S_+^{k_0} S_-^{l_0} \Lambda_0$, where $m := k_0 + l_0 = -\mathtt{tb}(L_0)$ and $k_0 - l_0 = \mathtt{rot}(L_0)$.

The strategy for proving Theorem 4 is now parallel to [1]. First determine an upper bound $\overline{\mathtt{tb}}(L_1)$ on the Thurston–Bennequin invariant of Legendrian realisations L_1 of (p, q)–cables of the given L_0, and show that Legendrian realisations with non-maximal `tb` destabilise. Show further that Legendrian realisations with maximal `tb` are determined by `rot`. If there are distinct Legendrian realisations with maximal `tb`, one also needs to understand the relationship between their stabilisations. This last issue will only arise in one subcase, where it can essentially be settled by a reference to [1].

By choosing the orientation of L_1 appropriately, we may assume $q \geq 0$ (and $p = 1$ for $q = 0$).

Case 1: $p = 0$ and $q = 1$. This is the case dealt with in Theorem 2.

Case 2: $p = 1$ and $q = 0$. Here L_1 is a meridian of L_0 and thus in particular a trivial knot. So the Thurston–Bennequin inequality gives $\overline{\mathtt{tb}}(L_1) = -1$, and this upper bound is obviously realised by a standard Legendrian unknot linked once with L_0.

By Theorem 3 we may assume that L_1 lies on a convex **standardly embedded** torus T, i.e. a torus isotopic in $J^1(S^1) \setminus L_0$ to a standard torus containing L_0 in the interior. Choose $R > 0$ large enough such that the standard torus $T_\infty := \partial M_R$ of slope ∞ contains T in the interior. Since $\mathtt{tb}(L_0) = -m$, we find a small tubular neighbourhood of L_0, contained in the interior of T, with convex boundary having two dividing curves of slope $-1/m$. Since slopes of convex standardly embedded tori decrease as we move away from L_0, and the slope can never be zero by the tightness of ξ, we see that the slope of T must be of the form $-r/s$ with coprime $r \in \mathbb{N}$, $s \in \mathbb{N}_0$, and $-r/s \leq -1/m$. By Giroux flexibility, cf. [6, Thm. 3.4], we may perturb T_∞ so that it still has two dividing curves of slope ∞, but now a Legendrian ruling of slope 0.

Let A be an annulus connecting L_1 and a Legendrian ruling curve L_∞ on T_∞. We have $t_A(L_1) = t_T(L_1)$ and $t_A(L_\infty) = t_{T_\infty}(L_\infty)$, so by Kanda's theorem — and using the convexity of both T and T_∞ — we may assume that A is convex. If $2n$ denotes the number of dividing curves on T, the algebraic intersection number of L_1 with the dividing set Γ_T on T equals $2nr$. Thus, if $n > 1$ or $r > 1$, there is a boundary parallel dividing curve on A along L_1, which allows us to destabilise that knot (obviously this can only happen if $\mathtt{tb}(L_1)$ was smaller than -1 to start with). If $n = r = 1$, but the geometric intersection number between L_1 and Γ_T is larger than the algebraic intersection number, we can again destabilise. If $n = r = 1$ and $\#(L_1 \cap \Gamma_T) = 2$, then $\mathtt{tb}(L_1) = t_A(L_1) = t_T(L_1) = -1$ by Theorem 3, so we are in the case $\mathtt{tb}(L_1) = \overline{\mathtt{tb}}(L_1)$. The dividing set of A then consists of two curves connecting L_1 with L_∞, so by Giroux flexibility we may assume that A is foliated

by Legendrian circles parallel to the boundary. This Legendrian ruling defines a Legendrian isotopy between L_1 and L_∞ (in the complement of L_0), which proves uniqueness of the Legendrian link $\mathbb{L}$.

Case 3: $p \geq 1$ and $q \geq 1$. Given a Legendrian link in $J^1(S^1)$, we are going to study it via its image in $S^3 \setminus K_0$ under the contactomorphism f mentioned in Section 2. We label objects in S^3 with a prime, and objects in $J^1(S^1)$ without. It will be convenient (for comparison with the classification in [1]) to define $L_0' := f(L_0)$ and $L_1' := \overline{f(L_1)}$ (i.e. $f(L_1)$ with reversed orientation). It is a straightforward check that if $\mathbb{L} = L_0 \sqcup L_1$ is a (p,q)–cable link in $J^1(S^1)$, then $\mathbb{L}' := L_0' \sqcup L_1'$ is a (p',q')–cable link in S^3, where $p' = -q$ and $q' = q - p$. (Here meridian μ' and longitude λ' on a torus T' around L_0' are chosen as in Figure 3.) With $m := -\mathtt{tb}(L_0)$ and $m' := -\mathtt{tb}(L_0')$ we have $-m = -m' + 1^2 = -m' + 1$. (See [1, p. 138] and the proof of Proposition 5 below for the relation between the Thurston–Bennequin invariant in $J^1(S^1)$ and that same invariant in S^3.)

However, we shall see that the classification of Legendrian cable links in S^3 does not translate directly into a classification of Legendrian cable links in $J^1(S^1)$.

Proposition 5. *If $p \geq 1$ and $q \geq 1$, then $\overline{\mathtt{tb}}(L_1) = p(q-1)$.*

Proof. In Figure 4 we exhibit the front projection of a (p,q)–cable L_1 to Λ_0 (or any of its stabilisations) with $p, q \geq 1$ and $\mathtt{tb}(L_1) = p(q-1)$ (which can be read off as the writhe of that front). This example shows that $\overline{\mathtt{tb}}(L_1) \geq p(q-1)$.



FIGURE 4. (p,q)–cable with maximal $\mathtt{tb}$ in Case 3.

It remains to prove the complementary inequality $\mathtt{tb}(L_1) \leq p(q-1)$ for every Legendrian (p,q)–cable L_1 of L_0.

If $p \geq q$, then the $(-p',-q')$–torus knot $f(L_1)$ is a *positive* torus knot. The maximal Thurston–Bennequin invariant of a positive $(-p',-q')$–torus knot is, according to [3, Thm. 4.1], given by $p'q' + p' + q'$ (this also holds for $q' = 0$, in which case $p' = -1$). With the correction term q^2 — for a knot homotopic to q times the generator of $\pi_1(J^1(S^1))$ — between the Thurston–Bennequin invariant in $J^1(S^1)$ and that in S^3 (cf. [1, p. 138]), we find

$$\mathtt{tb}(L_1) \leq (p-q)q - (p-q) - q + q^2 = p(q-1).$$

If $0 < p < q$, we consider the $(-q',-p')$–cable link $K_0 \sqcup f(L_1)$; here $(-q',-p') = (p-q,q)$ with $q \geq 2$. Thus, we produce a cable link in S^3 corresponding to Case 3(b) of [1]. Then [1, Lemma 4.6] tells us that $\mathtt{tb}(f(L_1)) \leq p'q'$. (Observe that the situation discussed there, where we might have to interchange the roles of the

two link components, does not arise: this is a consequence of $\mathtt{tb}(K_0) = -1 > -q$; see the discussion around [1, Lemma 4.5].) We compute

$$\mathtt{tb}(L_1) = \mathtt{tb}(f(L_1)) + q^2 \leq -q(q-p) + q^2 = pq.$$

So in all cases $p, q \geq 1$ we have at least the estimate $\mathtt{tb}(L_1) \leq pq$, and the following argument applies. (As regards the proof of the proposition, this argument is of course superfluous if $p \geq q$, but later it will be employed to conclude the classification in Case 3, where we need it for all $p, q \geq 1$.) Let T be a standardly embedded torus on which L_1 sits. Then $t_T(L_1) = \mathtt{tb}(L_1) - pq \leq 0$, so Theorem 3 allows us to assume that T is convex. As in Case 2, the dividing set Γ_T consists of $2n$ curves of slope $-r/s \leq -1/m$. Thus

$$\#(L_1 \cap \Gamma_T) \geq 2n \left| \begin{array}{cc} p & -s \\ q & r \end{array} \right| \geq 2p.$$

Hence $t_T(L_1) \leq -p$ (again by Theorem 3), and therefore $\mathtt{tb}(L_1) = pq + t_T(L_1) \leq p(q-1)$, as claimed. □

Remark. For $0 < p < q$, the estimate $\mathtt{tb}(L_1) \leq p(q-1)$ can also be obtained by appealing directly to [1, Lemma 4.6], applied to the $(p-q, q)$–cable link $K_0 \sqcup f(L_1)$. Since $\mathtt{tb}(K_0) = -1$, that lemma yields

$$\mathtt{tb}(L_1) = \mathtt{tb}(f(L_1)) + q^2 \leq (p-q)q - \max(1 \cdot (p-q) + q, 0) + q^2 = p(q-1).$$

Now the reasoning for Case 3 is entirely analogous to Case 2. We take T_∞ to be a (sufficiently large) convex standardly embedded torus having two dividing curves of slope ∞, and $L_\infty \subset T_\infty$ a ruling curve in a Legendrian ruling of slope q/p. Then $\#(L_\infty \cap \Gamma_{T_\infty}) = 2p$.

On the other hand, we have $\#(L_1 \cap \Gamma_T) \geq 2p$, with equality if and only if $r = 1$, $s = 0$, $n = 1$, and geometric intersection number being equal to the algebraic one. In all other cases, we can destabilise L_1. If $\mathtt{tb}(L_1) = \overline{\mathtt{tb}}(L_1)$, we must have $\#(L_1 \cap \Gamma_T) = 2p$, and with the help of a convex annulus A between L_1 and L_∞ we show those two curves to be Legendrian isotopic as in Case 2. This concludes the proof of Theorem 4 in Case 3.

The next corollary shows that in some cases a Legendrian (p', q')–cable link in $S^3 \setminus K_0$ can be further destabilised in S^3, but not in $S^3 \setminus K_0$.

Corollary 6. *For $0 < p < q-1$ there are Legendrian (p', q')–cable links in S^3 that are not Legendrian isotopic to the image under f of a Legendrian (p, q)–cable link in $J^1(S^1)$.*

Proof. Let $L_0' \sqcup L_1'$ be a Legendrian (p', q')–cable link in S^3 with $q' \geq 2$, $p' < 0$, and the unknotted component L_0' having $\mathtt{tb}(L_0') = -m'$. According to [1, Lemma 4.6], the maximal Thurston–Bennequin invariant $\overline{\mathtt{tb}}(L_1')$ is given by $p'q' - \max(m'p' + q', 0)$. If $L_0' \sqcup L_1' = f(L_0) \sqcup \overline{f(L_1)}$ is the image of a Legendrian (p, q)–cable link in $J^1(S^1)$ with $\mathtt{tb}(L_0) = -m$, then

$$m'p' + q' = (m+1) \cdot (-q) + q - p = -mq - p < 0.$$

If we consider arbitrary Legendrian realisations $L_0^* \sqcup L_1^*$ of this link type in S^3 (with $\mathtt{tb}(L_0^*) = -m'$), not necessarily coming from a link in $J^1(S^1)$, then $\overline{\mathtt{tb}}(L_1^*) = p'q' = pq - q^2$. This is larger than

$$\overline{\mathtt{tb}}(f(L_1)) = \overline{\mathtt{tb}}(L_1) - q^2 = p(q-1) - q^2,$$

so a sufficiently high destabilisation of L_1' in $S^3\setminus L_0'$ gives a Legendrian (p',q')–cable link which is not Legendrian isotopic to the image of a Legendrian (p,q)–cable link in $J^1(S^1)$. □

Thus, if $L_0\sqcup L_1$ is a Legendrian (p,q)–cable link in $J^1(S^1)$ with $0<p<q-1$ and $\mathtt{tb}(L_1)=\overline{\mathtt{tb}}(L_1)=p(q-1)$, then $\overline{f(L_1)}$ can be destabilised in $S^3\setminus f(L_0)$, but not in $S^3\setminus(f(L_0)\cup K_0)$. Still, our uniqueness proof above implies that there can be no choice for $\mathtt{rot}(L_1)$. This may seem surprising, since the classification from [1], applied to $f(L_0)\sqcup\overline{f(L_1)}$ in S^3, allows freedom in $\mathtt{rot}(\overline{f(L_1)})$ if $\mathtt{tb}(f(L_1))<\overline{\mathtt{tb}}(L_1')$. The paradox is resolved by considering, again, $K_0\sqcup f(L_1)$. Then [1, Lemma 4.11] gives

$$\mathtt{rot}(L_1)=\mathtt{rot}(f(L_1))=(p-q)\cdot\mathtt{rot}(K_0)=0.$$

This is indeed the rotation number realised by the cable in Figure 4.

Case 4: $p<0$ and $q\geq 1$. Here we have $p'=-q<0$ and $q'=q-p\geq 2$, so we are in Case 3(b) of [1]. Recall from the proof of Corollary 6 that $m'p'+q'=-mq-p$.

Case 4(a): $p<0$, $q\geq 1$ and $mq+p<0$. This puts us in Case 3(b1) of [1]. The Legendrian (p',q')–cable link $L_0'\sqcup L_1'$ in S^3 realising $\overline{\mathtt{tb}}(L_1')$, constructed explicitly in Section 5 of our earlier paper, sits in $S^3\setminus K_0$ and thus gives a Legendrian realisation of a (p,q)–cable link in $J^1(S^1)$ with

$$\mathtt{tb}(L_1)=\overline{\mathtt{tb}}(L_1)=p'q'-(m'p'+q')+q^2=pq+mq+p.$$

Here is a more direct description of such a Legendrian cable link realising $\overline{\mathtt{tb}}(L_1)$. Consider a small standard neighbourhood of L_0 with convex boundary T_0 having two dividing curves of slope $-1/m$. Since $q/p\neq -1/m$, Giroux flexibility allows us to assume that T_0 has a Legendrian ruling of slope q/p. Let L_1^0 be one of these ruling curves. Then, by Kanda's theorem,

$$\mathtt{tb}(L_1^0)=pq+t_{T_0}(L_1^0)=pq-\begin{vmatrix} p & m\\ q & -1\end{vmatrix}=pq+mq+p=\overline{\mathtt{tb}}(L_1).$$

The following argument is analogous to Cases 2 and 3, except that the protagonist is now T_0 in place of T_∞. Thus, let L_1 be any other Legendrian (p,q)–cable of L_0, lying on some standardly embedded torus T. The neighbourhood in the previous construction can be chosen arbitrarily small, so we may assume that T_0 sits in the interior of T. The fact that

$$t_T(L_1)=\mathtt{tb}(L_1)-pq\leq\overline{\mathtt{tb}}(L_1)-pq=mq+p<0$$

allows us to assume that T is convex, having $2n$ dividing curves of slope $-r/s\leq -1/m$.

Observe that, since $q>0$, we have $sq+rp\leq r(mq+p)<0$. Hence

$$\#(L_1\cap\Gamma_T)\geq -2\begin{vmatrix} p & -s\\ q & r\end{vmatrix}\geq -2(mq+p)=\#(L_1^0\cap\Gamma_{T_0}),$$

with equality if and only if Γ_T consists of two dividing curves of slope $-r/s=-1/m$. With the help of a convex annulus between L_1^0 and L_1 we conclude that L_1 can be destabilised for $\mathtt{tb}(L_1)<\overline{\mathtt{tb}}(L_1)$, and that L_1 is Legendrian isotopic to L_1^0 if its $\mathtt{tb}$ is maximal.

Case 4(b): $p<0$, $q\geq 1$ and $mq+p\geq 0$. This corresponds to Case 3(b2) of [1]. If $q=1$, we are in subcase 3(b2-iii); if $q>1$, in subcase 3(b2-ii). Here we

have $\overline{\mathtt{tb}}(L_1') = p'q' = pq - q^2$, hence $\mathtt{tb}(L_1) \leq pq$. There are several Legendrian realisations of this maximal $\mathtt{tb}$, distinguished by $\mathtt{rot}(L_1')$. All these realisations, as described in [1, Section 5], live in $S^3 \setminus K_0$ in such a way that we may regard them as links in $J^1(S^1)$. This yields $\overline{\mathtt{tb}}(L_1) = pq$. In order to establish a one-to-one correspondence between Legendrian (p,q)–cable links in $J^1(S^1)$ and (p',q')–cable links in S^3, we need to show the following:

- In a (p',q')–cable link in $S^3 \setminus K_0$ with non-maximal $\mathtt{tb}(L_1')$, the component L_1' can be destabilised inside $S^3 \setminus K_0$.
- Two (p',q')–cable links in $S^3 \setminus K_0$ with maximal $\mathtt{tb}(L_1')$ and the same classical invariants are Legendrian isotopic inside $S^3 \setminus K_0$.
- The relation between stabilisations with the same classical invariants coming from links with $\mathtt{tb}(L_1') = \overline{\mathtt{tb}}(L_1')$ but different rotation numbers are as described in Lemmata 4.20 and 4.21 of [1], respectively.

The first point we can settle directly in $J^1(S^1)$. If $L_0 \sqcup L_1$ is a (p,q)–cable link with $\mathtt{tb}(L_1) < pq$, we find a convex standardly embedded torus T on which L_1 sits. If the actual geometric intersection number between L_1 and the dividing set Γ of T is larger than the minimal one possible, there exists a bypass for L_1 on T, and we can destabilise L_1. Thus, we may assume that T is in standard form of slope $-r/s \leq -1/m$, and the non-maximality of $\mathtt{tb}(L_1)$ then forces $-r/s \neq q/p$. Further, the condition $mq + p \geq 0$ gives $q/p \leq -1/m$. So we find a convex torus $T_{q/p}$ of slope q/p either between T_0 and T (if $q/p > -r/s$) or between T and T_∞ (if $q/p < -r/s$), and we can use a convex annulus between L_1 and a Legendrian divide on $T_{q/p}$ to destabilise L_1.

The second point, for $q > 1$, requires the use of [1, Lemma 4.14]. The proof of that lemma goes through in $S^3 \setminus K_0$, as can be seen by thinking of S^3 as being composed of a union of four pieces: a standard neighbourhood of L_0 with boundary T_0, a thickened torus between T_0 and T (on which L_1 sits), a thickened torus between T and T_∞, and a standard neighbourhood of K_0.

For $q = 1$, this point cannot be settled with a simple reference to [1], for this is exactly the situation that necessitated the discussion in Section 4. That argument allows us to assume that our cable link sits in the interior of a solid torus V with convex boundary having two dividing curves of slope $1/p$. (The reduction of the number of dividing curves on ∂V to 2 by using meridional discs in the complement also works in $S^3 \setminus K_0 = J^1(S^1)$, because we can always choose the bypasses in such a way that we do not move across K_0 in order to effect the destabilisation.)

By 'unwinding' this solid torus V as in Section 4, we may identify $L_0 \sqcup L_1$ with a link $L_0^* \sqcup L_1^*$ in $J^1(S^1)$ topologically isotopic to $\Lambda_0 \sqcup \Lambda_1$, with $\mathtt{tb}(L_0^*) = -m - p$ and $\mathtt{tb}(L_1^*) = p \cdot 1 - p = 0$. Thus, if $m + p > 0$, then this link is determined by its classical invariants even inside V. If $m + p = 0$, then $L_0^* \sqcup L_1^*$ is Legendrian isotopic to $\Lambda_0 \sqcup \Lambda_1$ or $\Lambda_1 \sqcup \Lambda_0$, and in particular a 2–copy. So $L_0 \sqcup L_1$ is likewise a 2–copy, but now of a stabilised knot. In that case, it is easy to see that permutation is possible by a Legendrian isotopy, see [8, Prop. 4.2a].

Finally, for the third point, a close inspection of the relevant arguments in [1] reveals that the mentioned lemmata do indeed hold in $S^3 \setminus K_0$.

This completes the discussion of Case 4 and hence the proof of Theorem 4.

Acknowledgements. Most of this research was carried out during a stay of F. D. at the Mathematical Institute of the Universität zu Köln, supported by DAAD grant A/06/27941. F. D. is partially supported by grant no. 10631060 of the National Natural Science Foundation of China.

H. G. would like to thank the American Institute of Mathematics for its support during the workshop "Legendrian and transverse knots" (September 2008), where the final writing was done, and the organisers D. Fuchs, S. Tabachnikov and L. Traynor for the invitation to that workshop.

References

[1] F. Ding and H. Geiges, Legendrian knots and links classified by classical invariants, *Commun. Contemp. Math.* **9** (2007), 135–162.
[2] J. B. Etnyre, Legendrian and transversal knots, in *Handbook of Knot Theory*, eds. W. Menasco and M. Thistlethwaite (Elsevier, Amsterdam, 2005), pp. 105–185.
[3] J. B. Etnyre and K. Honda, Knots and contact geometry I: Torus knots and the figure eight knot, *J. Symplectic Geom.* **1** (2001), 63–120.
[4] J. B. Etnyre and K. Honda, On the nonexistence of tight contact structures, *Ann. of Math. (2)* **153** (2001), 749–766.
[5] J. B. Etnyre and K. Honda, On connected sums and Legendrian knots, *Adv. Math.* **179** (2003), 59–74.
[6] K. Honda, On the classification of tight contact structures I, *Geom. Topol.* **4** (2000), 309–386.
[7] Y. Kanda, On the Thurston–Bennequin invariant of Legendrian knots and nonexactness of Bennequin's inequality, *Invent. Math.* **133** (1998), 227–242.
[8] K. Mishachev, The N–copy of a topologically trivial Legendrian knot, *J. Symplectic Geom.* **1** (2003), 659–682.
[9] L. Ng and L. Traynor, Legendrian solid-torus links, *J. Symplectic Geom.* **2** (2004), 411–443.
[10] L. Traynor, Legendrian circular helix links, *Math. Proc. Cambridge Philos. Soc.* **122** (1997), 301–314.

Department of Mathematics, Peking University, Beijing 100871, P. R. China
E-mail address: dingfan@math.pku.edu.cn

Mathematisches Institut, Universität zu Köln, Weyertal 86–90, 50931 Köln, Germany
E-mail address: geiges@math.uni-koeln.de